

See discussions, stats, and author profiles for this publication at:
<https://www.researchgate.net/publication/301576740>

EPINEPHRINE;A VASOCONSTRICTOR OR A UTERINE RELAXANT? A CASE SERIES

Article · April 2016

DOI: [10.17957/TPMJ/16.3297](https://doi.org/10.17957/TPMJ/16.3297)

READS

6

5 authors, including:



Farnoush Farzi

14 PUBLICATIONS 11 CITATIONS

[SEE PROFILE](#)



Azadeh Raoufi

Mehr medical institute

7 PUBLICATIONS 9 CITATIONS

[SEE PROFILE](#)

All in-text references underlined in blue are linked to publications on ResearchGate, Available from: Azadeh Raoufi letting you access and read them immediately. Retrieved on: 05 July 2016



Associate Professor of
Anesthesiology, Anesthesia
Research Center, School of
Medicine, Guilan University of
Medical Sciences, Rasht, Iran
3. M.Sc of Developmental Biology, Anesthesia
Research Center
4. M.D Ali Mirmansouri¹, Farnoush
Farzi², Azadeh Raoufi³, Ziba Zahiri
Sorouri⁴,

Professor of Obstetric and Fereidon Mortazavi Najafabadi⁵ Gynecology, Reproductive

Health Research Center, Alzahra **ABSTRACT:** The most common complication of spinal anesthesia for cesarean section is Hospital, School of Medicine, Guilan hypotension. Ephedrine is the most commonly used vasopressor that increases blood pressure University of Medical Sciences, with minimal impact on uteroplacental blood flow. An alpha-1 adrenergic receptor agonist may Rasht, Iran need to be administrated when ephedrine is ineffective. Unavailability of alpha-1 receptor 5 . M.D Assistant Professor of agonists in a period of time in our center leads to administration of epinephrine as the second

Anesthesiology drug. In the present study, the data of 14 patients with ephedrine resistant hypotension during spinal anesthesia for cesarean section were reviewed. Increase in maternal blood pressure **Correspondence:**

Farnoush farzi M.D was recorded one minute after epinephrine administration in all patients. Surprisingly, this Anesthesia Research Center, Valayt medication also causes uterine relaxation after one to five minutes. Ease of fetal extraction Hospital, School of Medicine, Guilan was noticed in 13 patients. All patients achieved adequate uterine contraction after delivery.

University of Medical Sciences, Rasht,

Epinephrine helped regulate blood pressure and surprisingly facilitate uterine relaxation in

Iran

Farnoush_farzi@gums.ac.ir
farnoush_farzi@yahoo.com

patients with emergency cesarean section with spinal anesthesia.

Key words:

Hypotension; Epinephrine; Anesthesia, Spinal; Cesarean Section; Tocolytic

Article received on: 18/02/2016

Accepted for publication: Article Citation: Mirmansouri A, Farzi F, Raoufi A, Zahiri Sorouri Z, Mortazavi Najafabadi 10/03/2016 F. Epinephrine; a vasoconstrictor or a uterine relaxant? a case series.

Received after proof reading:

12/04/2016

Professional Med J 2016;23(4):504-508. DOI: 10.17957 /TPMJ/ 16.3297

INTRODUCTION

Spinal anesthesia has become preferred choice for anesthesia in emergency cesarean delivery. In obstetrics, regional anesthesia particularly spinal block, eliminate the need for general anesthesia. However, hypotension is commonly found after performing of spinal anesthesia and continues hypotension can have detrimental maternal and fetal effects.^{1,2} Therefore it is crucial to prevent and treat induced hypotension. Many strategies have been developed to minimize the impact of hypotension but if occurs, use of vasopressors are needed.³

Ephedrine that its action is considered to be indirect is the most commonly used vasopressor. Repeated doses of ephedrine may be needed to induce maternal dynamic stability but it should be noted that tachyphylaxis can occur following repeated use of this drug.⁴ So an alpha-1 adrenergic receptor agonist may require to be added when ephedrine is inefficient. Epinephrine is not ordinary used as an agent to treat spinal hypotension. But in the present study because

of restriction on access to other drugs and the availability of epinephrine, therefore it is reasonable for epinephrine administration to treat hypotension. As well, epinephrine acts as an agonist for smooth muscle beta2-adrenoceptors activation.⁵ Beta2-adrenoceptors are found abundantly in the smooth muscle of uterus that mediating relaxation.⁶

In this case series, we present 14 patients with ephedrine resistant hypotension during spinal anesthesia for emergency cesarean delivery that all of them were treated with low dose of epinephrine.

CASE REPORT

In this case series, the medical records of patients with emergency cesarean delivery at Alzahra hospital between 2007 and 2011, who underwent spinal anesthesia, were reviewed. During study period, 14 patients with ephedrine resistant hypotension were identified. The mean age of patients was 25.3 years (range from 16 to 35). Demographic and anesthetic data are

summarized in table-I and II. Neonatal outcome are shown in table-III.

Briefly, on arrival at the operating room, basal monitoring (including non-invasive blood pressure measurement, heart rate, respiratory rate, arterial oxygen saturation) were started. Sixteen or eighteen gauge intravenous cannula was used for administration of normal saline (5 cc/kg).

Spinal anesthesia was performed with hyperbaric 5% lidocaine (70 mg) in the sitting position with 25 G Quincke needle. Patients were then immediately returned to the supine position. Systolic blood pressure was measured at 2-3 min intervals. If blood pressure fell below 100 mmHg, ephedrine (10mg) was administrated intravenously. Further ephedrine blouses (10 mg) were given when systolic blood pressure remains low. Epinephrine (10 μ g) was administered for treating of ephedrine resistant hypotension. All patients received intravenous oxytocin (30IU) infusion over 1 hour period. If uterine atony persisted after the administration of oxytocin, methylergonovine (0.2 mg) was administrated intramuscularly. Anesthesiologist provided neonatal

resuscitation when it was required. After delivery, apgar scores were determined at 1th, 5th and 10thmin. Repeated dose of ephedrine displayed tachyphylaxis in all patients so epinephrine were given as the second drug for treating hypotension induced by spinal anesthesia. Increase in maternal blood pressure was recorded 1 min after epinephrine administration. Surprisingly, this medication also causes uterine relaxation in all patients after 1 to 5 min. Ease fetal extraction was indicated in thirteen patients. Neonates were delivered in case of multiple gestations, dystocia and fetal distress, had low apgar scores at first minute and after resuscitation, apgar scores increase at fifth minute after birth.

Three of fourteen patients required more than 30 unit of oxytocin infusion to achieve uterine contraction. All patients achieved adequate uterine contraction after delivery. Methylergonovine was given to four patients (including previous cesarean, multiple gestation, breech presentation and IUFD) because of uterine atony persists after the administration of oxytocin.

There was no sign of massive uterine bleeding and blood transfusion wasn't required. There were no

| Patient | Age (yrs) | Weight (kg) | Obstetrics records | | | | | Gestational age(weeks) | Fetal presentation | Reason for cesarean | Current medication |
|---------|-----------|-------------|--------------------|---|----|---|---|------------------------|--------------------|--|-----------------------|
| | | | D | L | Ab | P | G | | | | |
| 1 | 34 | 79 | - | - | - | 1 | 1 | 34 | Breech | Premature rupture of membranes | - |
| 2 | 31 | 75 | - | 1 | - | 2 | 2 | 39 | Cephalic | Previous cesarean | |
| 3 | 35 | 96 | 0 | 0 | 0 | 0 | 0 | 37 | Cephalic | Fetal distress | Insulin |
| 4 | 17 | 75 | 0 | 0 | 0 | 1 | 1 | 40 | Cephalic | Dystocia | - |
| 5 | 22 | 70 | 0 | 0 | 0 | 0 | 1 | 28 | Cephalic | Multiple gestation | - |
| 6 | 16 | 50 | 0 | 0 | 0 | 0 | 1 | 36 | Breech | Premature rupture | - |
| 7 | 18 | 60 | 0 | 0 | 0 | 0 | 1 | 39 | Cephalic | Fetal distress | - |
| 8 | 33 | 70 | 0 | 0 | 0 | 0 | 1 | 30 | Transverse | Multiple gestation-Prematurity-preterm Labor | Levothyroxine |
| 9 | 27 | 60 | 0 | 0 | 0 | 0 | 1 | 34 | Breech | Oligohydramnios | - |
| 10 | 23 | 68 | 0 | 0 | 0 | 1 | 1 | 26 | Breech | Footling | Levothyroxine |
| 11 | 23 | 67 | 0 | 1 | 2 | 2 | 4 | 38 | Breech | Previous cesarean | - |
| 12 | 19 | 65 | 0 | 0 | 0 | 0 | 1 | 41 | Cephalic | Fetal distress | |
| 13 | 22 | 60 | 0 | 0 | 0 | 2 | 2 | 34 | Cephalic | Previous cesarean | Smoking and addiction |
| 14 | 34 | 80 | 0 | 1 | 0 | 1 | 2 | 37 | Cephalic | IUFD | |

Table-I. Demographic data of 14 patients with ephedrine resistant hypotension during spinal anesthesia for emergency cesarean delivery.

D, dead; L, live; Ab, abortion; P, para; G, gravid; IUFD, intrauterine fetal death

cases of emergency hysterectomy due to persistent uterine atony unresponsive to medication.

| Patient | Preoperative | | | induction | | | | | | | | | | | | | | |
|---------|--------------|---------|----------|-----------------------------------|----------|-----------------------------------|----------|-----------------------------------|----------|------------------------------------|----------|---|----------|--|----------|--|----------|---------|
| | | | | 1 st min of anesthesia | | 3 rd min of anesthesia | | 5 th min of anesthesia | | 10 th min of anesthesia | | Immediately before epinephrine administration | | 1 st min after epinephrine administration | | 5 th min after epinephrine administration | | |
| | BP(mmHg) | HR(bpm) | BP(mmHg) | HR(bpm) | BP(mmHg) | HR(bpm) | BP(mmHg) | HR(bpm) | BP(mmHg) | HR(bpm) | BP(mmHg) | HR(bpm) | BP(mmHg) | HR(bpm) | BP(mmHg) | HR(bpm) | BP(mmHg) | HR(bpm) |
| 1 | 120/80 | 60 | 90/60 | 65 | 80/60 | 60 | 90/60 | 70 | 90/60 | 75 | 85/60 | 90 | 100/70 | 100 | 120/70 | 120 | | |
| 2 | 120/80 | 70 | 100/70 | 80 | 90/60 | 80 | 90/60 | 80 | 90/60 | 80 | 90/60 | 80 | 100/70 | 90 | 110/70 | 95 | | |
| 3 | 110/70 | 90 | 90/60 | 85 | 80/60 | 85 | 80/60 | 85 | 100/70 | 92 | 90/60 | 90 | 110/70 | 100 | 100/65 | 94 | | |
| 4 | 130/70 | 90 | 120/70 | 80 | 110/70 | 80 | 100/70 | 75 | 90/60 | 75 | 90/60 | 75 | 110/70 | 85 | 140/80 | 90 | | |
| 5 | 120/75 | 130 | 110/70 | 120 | 85/65 | 120 | 95/75 | 115 | 90/80 | 110 | 80/70 | 120 | 110/85 | 130 | 100/80 | 120 | | |
| 6 | 110/80 | 80 | 70/50 | 90 | 90/60 | 90 | 80/60 | 95 | 100/75 | 90 | 80/60 | 95 | 110/80 | 112 | 100/75 | 110 | | |
| 7 | 120/80 | 70 | 110/80 | 80 | 95/80 | 85 | 110/75 | 80 | 110/75 | 75 | 90/65 | 80 | 120/80 | 85 | 125/80 | 100 | | |
| 8 | 120/80 | 80 | 100/70 | 80 | 100/70 | 80 | 100/70 | 85 | 100/70 | 85 | 85/60 | 85 | 110/70 | 90 | 120/70 | 90 | | |
| 9 | 110/70 | 70 | 100/70 | 75 | 100/70 | 75 | 100/70 | 75 | 100/70 | 75 | 90/65 | 75 | 110/70 | 85 | 110/70 | 90 | | |
| 10 | 110/70 | 90 | 100/70 | 95 | 100/70 | 95 | 100/70 | 95 | 100/70 | 95 | 90/65 | 95 | 106/70 | 105 | 105/70 | 105 | | |
| 11 | 110/70 | 70 | 100/70 | 60 | 100/70 | 70 | 120/80 | 75 | 120/80 | 75 | 95/65 | 70 | 120/80 | 75 | 120/80 | 80 | | |
| 12 | 120/80 | 70 | 100/70 | 100 | 100/70 | 100 | 90/70 | 105 | 90/70 | 105 | 85/50 | 100 | 110/80 | 110 | 115/80 | 105 | | |
| 13 | 120/80 | 70 | 100/70 | 75 | 90/70 | 80 | 100/70 | 75 | 95/75 | 80 | 85/55 | 70 | 120/80 | 80 | 140/80 | 90 | | |
| 14 | 100/60 | 95 | 90/50 | 90 | 100/50 | 85 | 100/60 | 85 | 110/60 | 85 | 90/65 | 85 | 110/50 | 95 | 110/60 | 95 | | |

Table-II. Anesthetic data of 14 patients with ephedrine resistant hypotension during spinal anesthesia for emergency cesarean delivery.

BP, blood pressure; HR, heart rate

| Patient | Onset of tocolysis after epinephrine administration(min) | Fetal extraction difficult, normal,easy | Apgar score at 1 min | Apgar score at 5 min | Apgar score at 10 min | Total dose of oxytocin IU (infusion) | Administered methylergonovine mg | Uterine Contraction after delivery |
|---------|--|---|----------------------|----------------------|-----------------------|--------------------------------------|----------------------------------|------------------------------------|
| 1 | 2 | Easy | 8 | 9 | 9 | 30 | | good |
| 2 | 2 | Easy | 1 | 7 | 8 | 30 | 0.2 | good |
| 3 | 1 | Easy | (5 | 8 | 8 | 30 | | good |
| 4 | 4 | easy | 5 | 7 | 8 | 30 | | good |
| 5 | 3 | easy | 5-5-5-4 | 7-7-7-6 | 7-7-7-7 | 30 | 0.2 | good |
| 6 | 2 | easy | 7 | 8 | 9 | 30 | | good |
| 7 | 1 | normal | 7 | 9 | 9 | 30 | | good |
| 8 | 2 | easy | 2-5-5 | 5-7-7 | 6-8-8 | 50 | | good |

| | | | | | | | | |
|----|---|-----------|---|---|---|----|-----|------|
| 9 | 1 | easy | 7 | 8 | 9 | 50 | 0.2 | good |
| 10 | 1 | easy | 5 | 7 | 7 | 30 | | good |
| 11 | 5 | difficult | 7 | 8 | 8 | 30 | | good |
| 12 | 5 | normal | 4 | 6 | 8 | 30 | | good |
| 13 | 3 | easy | 4 | 8 | 8 | 30 | | good |
| 14 | 2 | normal | 0 | 0 | 0 | 40 | 0.2 | Good |

Table-III. Neonatal outcome of 14 patients with ephedrine resistant hypotension during spinal anesthesia for emergency cesarean delivery.

Professional Med J 2016;23(4): 504-508.

www.theprofesional.com

506

DISCUSSION

The findings of the present case series include that epinephrine helped regulate blood pressure and also induced uterine relaxation in patients with emergency cesarean section with spinal anesthesia. Surgeon admitted ease of fetal extraction and no patients experienced unusual severe bleeding, resistant uterine atony and emergency hysterectomy.

Spinal anesthesia is an effective technique whereas preventing general anesthesia and related risks in cesarean section. However, spinal anesthesia for cesarean section is not without risk, because of induced maternal hypotension which can cause maternal detrimental effects and fetal distress. Spinal anesthesia-induced hypotension is triggered by sympathetic blockade which leads to reduction in peripheral vascular resistance and venous return.⁷ Many strategies have been developed to minimize the impact of hypotension. In the present case series, we used ephedrine as the first line vasopressor for management of maternal spinal hypotension but after three 10 mg doses, tachyphylaxis was induced. Epinephrine 5-10 µg was injected intravenously when ephedrine was ineffective. Epinephrine isn't the preferred vasopressor for treating hypotension during spinal anesthesia for cesarean section, but should be available in operating room. In the present study, because of limited access to an alpha-1 adrenergic receptor agonist, in a period of time, epinephrine was used as the second line vasopressor. Surprisingly, epinephrine resulted uterine relaxation within 1 to 5 min and fetal were extracted easily. Emergent uterine

relaxation is required for situations like internal podalic version, retained placenta, inverted uterus and breech delivery during cesarean section.⁸ Halogenated anesthetics were used traditionally for smooth muscle uterine relaxation in emergency situation, but it is required endotracheal intubation using rapid sequence induction of anesthesia and can be a very risky option. However, a way of avoiding general anesthesia and providing transient uterine relaxation for full stomach patients would be beneficial.

Nitroglycerin, as a short acting smooth muscle relaxant acts via increase in synthesis of cyclic guanosine monophosphate (cGMP) and induction cGMP-mediated dephosphorylation of myosin light chain.⁹ Mayer¹⁰ discusses a case of cesarean section for twin pregnancy at 30 weeks. After initiation of spinal anesthesia, systolic blood pressure decreased from 120 to 95 mmHg. Following the request for uterine relaxation, nitroglycerin was administrated to facilitate fetal extraction. Systolic blood pressure also reduced to 85 mmHg but increased after ephedrine administration.

Despite tocolytic (uterine relaxant) effect, hypotensive properties of nitroglycerin during general anesthesia have been described.¹¹⁻¹³ As a result, an important concern about this drug in spinal anesthesia for cesarean section has been the excessive induced hypotension. So it can be assumed that nitroglycerin is not good selection for uterine relaxation in patients with hypotension due to sympathetic block after spinal anesthesia.

In general, drugs with myometrial beta-2 adrenoreceptor agonist activity have been used for uterine relaxation.¹⁴ Intravenous epinephrine, at low and high doses, can respectively stimulate beta 2, beta 1 and alpha 1 adrenoceptors. It relaxes the smooth muscles by activating beta-2 adrenoceptors and enhances the rate and force of heart contraction through beta-1 adrenoceptors.¹⁵ The combination of these effects of epinephrine is desirable during spinal

anesthesia for cesarean section. Beyond the effect on hypotension treatment, epinephrine is a surprisingly a cause of uterine relaxation in this study. The findings, generate hypothesis that epinephrine is a suitable choice, not only for regulation of blood pressure but also surprisingly for facilitation of uterine relaxation in patients with emergency cesarean section with spinal anesthesia induced hypotension needed uterine relaxation for fetal extraction delivery.

So in summary, the present case series provided information to introduce epinephrine, for treatment of ephedrine resistant hypotension and providing adequate uterine relaxation in spinal anesthesia for

[Professional Med J 2016;23\(4\): 504-508.](#)

cesarean section. However the findings should be expounded discreetly and considered as basic step that will inform subsequent studies on administration of epinephrine as a uterine relaxant. Until completion of studies, the usage of epinephrine should be avoided in hypertensive pregnant patients (preeclampsia and eclampsia). [Copyright© 10 Mar, 2016.](#)

REFERENCES

1. Rout CC, Rocke DA. [Prevention of hypotension following spinal anaesthesia for cesarean section.](#) Int Anesthesiol Clin. 1994;32(2):117-36.
2. Corke B, Datta S, Ostheimer G, Weiss J, Alper M. [Spinal anaesthesia for Caesarean section.](#) Anaesthesia. 1982;37(6):658-62.
3. Kee WDN, Khaw KS. [Vasopressors in obstetrics: what should we be using?](#) Curr Opin Anaesthesiol. 2006;19(3):238-43.

4. Persky AM, Berry NS, Pollack GM, Brouwer KL. [Modelling the cardiovascular effects of ephedrine.](#) Br J Clin Pharmacol. 2004;57(5):552-62.
5. Koike K, Ichino T, Horinouchi T, Takayanagi I. [The beta2-and beta3-adrenoceptor-mediated relaxation induced by isoprenaline and salbutamol in guinea pig taenia caecum.](#) J Smooth Muscle Res. 1997;33(3):99106.
6. Tanaka Y, Horinouchi T, Koike K. [New insights into beta-adrenoceptors in smooth muscle: distribution of receptor subtypes and molecular mechanisms triggering muscle relaxation.](#) Clin Exp Pharmacol Physiol. 2005;32(7):503-14.
7. Morgan PJ, Halpern SH, Tarshis J. [The effects of an increase of central blood volume before spinal anaesthesia for cesarean delivery: a qualitative systematic review.](#) Anesth Analg. 2001;92(4):9971005.
8. Dufour P, Vinatier D, Puech F. [The use of intravenous nitroglycerin for cervico-uterine relaxation: a review of the literature.](#) Arch Gynecol Obstet. 1997;261(1):1-7.
9. Murad F. [Cyclic guanosine monophosphate as a mediator of vasodilation.](#) J Clin Invest. 1986;78(1):1.
10. Mayer DC, Weeks SK. [Antepartum uterine relaxation with nitroglycerin at caesarean delivery.](#) Can J Anaesth. 1992;39(2):166-9.



[www.theprofesional.com](#)

507

11. Smith GN, Brien JF. [Use of nitroglycerin for uterine relaxation.](#) Obstet Gynecol Surv. 1998;53(9):559-65.
12. Axemo P, Fu X, Lindberg B, Ulmsten U, Wessen A. [Intravenous nitroglycerin for rapid uterine relaxation.](#) Acta Obstet Gynecol Scand. 1998;77(1):50-3.
13. Fahmy NR. [Nitroglycerin as a hypotensive drug during general anesthesia.](#) Anesthesiology. 1978;49(1):17-20.
14. Morrison JJ, Rennie JM. [Clinical, scientific and ethical aspects of fetal and neonatal care at extremely preterm periods of gestation.](#) Br J Obstet Gynaecol. 1997;104(12):1341-50.
15. Kaumann A, Hall J, Murray K, Wells F, Brown M. [A comparison of the effects of adrenaline and noradrenaline on human heart: the role of β1-and β2adrenoceptors in the stimulation of adenylate cyclase and contractile force.](#) Eur Heart J. 1989;10(suppl B):29-37.

AUTHORSHIP AND CONTRIBUTION DECLARATION

| Sr. # | Author's Full Name | Contribution to the paper | Author=s Signature |
|-------|--------------------|---------------------------|--------------------|
|-------|--------------------|---------------------------|--------------------|

| | | | |
|---|-----------------------------------|---|--|
| 1 | Ali Mirmansouri M.D | Study design, Data analysis |  |
| 2 | Farnoush Farzi M.D | Correspondence, Study design, Data analysis, data collection, |  |
| 3 | Azadeh Raoufi M.Sc | manuscript preparation Data |  |
| 4 | Ziba Zahiri Sorouri M.D | analysis, manuscript preparation Data |  |
| 5 | Fereidon Mortazavi Najafabadi M.D | collection Data collection |  |